



De novo onset or exacerbation of psoriasis following the SARS-CoV-2 infection or COVID vaccination | 1

Several studies reported that infection with the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and COVID-19 vaccination are associated with various cutaneous manifestations, either as flares of preexisting dermatoses or as *de novo* onset of immune-mediated dermatological disorders. Psoriasis is a chronic immune-mediated disease that can be exacerbated or manifest *de novo* following concomitant infections. In this case report, the authors from Italy presented a case series of bio-naïve patients who experienced exacerbations or *de novo* onsets of psoriasis following the SARS-CoV-2 infection or COVID-19 vaccination.

Psoriasis is an autoimmune disorder characterized by the hyperproliferation of epidermal keratinocytes with inflammation of the epidermis and dermis. It manifests with erythematous plaques covered with silvery scales, particularly over the extensor surfaces, scalp, and lumbosacral region. The exact pathophysiology is unknown. It is considered multifactorial, with environmental, genetic, and immunological factors involved. Trauma, infection, and certain medications are common triggers.



The authors noted that several reports have been published regarding psoriasis flares following COVID-19 vaccination or SARS-CoV-2 infection. However, no standardized guidelines have been proposed to address the treatment of these psoriasis cases specifically. According to a recent systematic review that analyzed seven studies reporting new-onset



De novo onset or exacerbation of psoriasis following the SARS-CoV-2 infection or COVID vaccination | 2

psoriasis, most flares developed following mRNA COVID-19 vaccination. Psoriasis was most commonly diagnosed following immunization with mRNA COVID-19 Pfizer/BioNTech vaccines. Most patients were treated with topical agents, while some patients with exacerbation of psoriasis received systemic treatment, including biologics.

About the Study and Results

The authors presented 28 patients who were bio-naïve and experienced flares or new onsets of psoriasis following COVID-19 infection or vaccination. Of these, 25 patients experienced an exacerbation, and three patients experienced a new onset of psoriasis. The mean age was 52.6 years, and 68% male.

Severe flares occurred in 25 patients (89%) diagnosed with mild psoriasis approximately 20 years before. Of these 25 patients, 64% (16 patients) experienced a flare after COVID-19 vaccination, and 36% (9 patients) after the SARS-CoV-2 infection.

Three patients were diagnosed with a new onset of the disease. One patient developed moderate plaque psoriasis after acute COVID-19. The other two patients experienced a new onset of psoriasis after receiving the mRNA Pfizer/BioNTech vaccines. The first patient developed moderate plaque psoriasis after the third dose of the Pfizer/BioNTech vaccine, while the second patient developed severe palmoplantar pustular and genital plaque psoriasis after the second dose of the Pfizer/BioNTech vaccine.



All patients received biological treatment according to the Italian guidelines for managing psoriasis. The evaluation of PASI (Psoriasis Area and Severity Index) at baseline and week 16 was used to determine the effectiveness of biological treatment. They all started taking an interleukin (IL) inhibitor after testing negative for viral hepatitis, HIV, and tuberculosis. Anti-IL-23 medications were the most frequently prescribed (20 patients), with risankizumab being the most chosen treatment of this category (12 patients), followed by guselkumab and tildrakizumab (4 patients each). Six patients received an IL-17 inhibitor (three secukinumab, one ixekizumab, one brodalumab, and one bimekizumab). Finally, two patients received ustekinumab, an anti-IL-12/23 drug. All patients were successfully managed, with higher rates of clinical responses after 16 weeks.

None of the patients receiving biological treatment experienced a flare of psoriasis, except for one female patient who developed a severe flare of generalized pustular psoriasis after receiving the second dose of the Pfizer/BioNTech vaccine and skipping the planned administration of ustekinumab. In this case, the immediate treatment with risankizumab led to almost complete skin clearance within 16 weeks. The authors stated that 12% of their patients receiving biological therapy reported mild worsening of plaque psoriasis following COVID-19 infection or vaccination.



Conclusion

This study described a case series of 28 patients who experienced *de novo* onset or flares of psoriasis following infection with SARS-CoV-2 or COVID-19 vaccination. All of these patients were successfully managed with anti-IL-17 or anti-IL-23 medications.

This article was published in the Journal of Dermatological Treatment.

Journal Reference

Gargiulo L, Ibba L, Vignoli CA, et al. New-onset and flares of psoriasis after COVID-19 infection or vaccination successfully treated with biologics: a case series, Journal of Dermatological Treatment, 2023, 34:1, 2198050 (Open Access)

<https://doi.org/10.1080/09546634.2023.2198050>